## **PEDIPHEN**

# Diphenhydramine Hydrochloride Syrup

#### DESCRIPTION

**PEDIPHEN** (Diphenhydramine Hydrochloride BP) is an antihistamine with anticholinergic and sedative effect.

### COMPOSITION

**PEDIPHEN** Syrup : Each 5 ml syrup contains Diphenhydramine Hydrochloride BP 10 mg.

#### **INDICATIONS**

- Seasonal, perennial and vasomotor rhinitis
- Urticaria, angioneurotic edema, anaphylaxis
- Pruritus
- Preanesthetic medication, emesis, motion sickness
- Miscellaneous include Meniere's disease and Parkinsonism

#### DOSAGE AND ADMINISTRATION

For Adults  $:2\frac{1}{2}$  to 5 teaspoonfuls 3 to 4 times daily, not exceed 300 mg in 24 hours. For Children: Under 2 years:  $\frac{1}{2}$  to  $\frac{1}{4}$  teaspoonful 1 to 3 times or as per physician advice. 2 to 6 years:  $\frac{1}{2}$  to 1 teaspoonful 1 to 3 times or as per physician advice. 6 to 12 years:  $1\frac{1}{4}$  to  $2\frac{1}{2}$  teaspoonfuls 3 to 4 times, not exceed 150 mg in 24 hours.

#### CONTRAINDICATION

Known hypersensitivity to Diphenhydramine Hydrochloride, CNS depression, patient taking monoamine oxidase (MAO) inhibitors. This drug should not be used in newborn or premature infants.

## PRECAUTION

If drowsiness, patient should not drive or operate machinery requiring alertness. Alcohol and certain other CNS depressants can potentiate the sedative effect of antihistaminics and patient should be warned of these effects.

#### SIDE EFFECTS

Drowsiness, dryness of mouth, sedation, disturbed coordination.

## PACKAGING

PEDIPHEN Syrup: Each bottle contains 100 ml syrup.

