Lodical Amlodipine

DESCRIPTION

Lodical (Amilodipine) is a calcium chanel antagonist of the Dihydropyridine group and inhibits the transmembrane influx of calcium ion into cardiac & smooth muscle.

Lodical 5 tablet: Each film coated tablet contains Amlodipine Besilate BP equivalent to Amlodipine 5 mg.

INDICATION

Hypertension: Lodical is indicated for the first line treatment of Hypertension. It may

be used alone or in combination with other antihypertensive agents.

Prophylaxis of Angina: Lodical is indicated for Prophylaxis of Angina. Lodical may be used alone or in combination with other antianginal agents.

 $\begin{tabular}{lll} \textbf{DOSAGE AND ADMINISTRATION} \\ \textbf{The usual oral dose in Hypertension or Angina is 5 mg once daily with a maximum} \\ \end{tabular}$ dose of 10 mg once daily.

CONTRAINDICATION

Amlodipine is contraindicated in patients with known hypersensitivity to Dihydropyridine (e.g. Amlodipine, Nifedipine, Nicardipine, Isradipine).

Since the vasodilatation induced by Amlodipine is gradual in onset, Acute Hypotension has rarely been reported after oral administration of Amlodipine. Nonetheless, caution should be exercised when administering Amlodipine with any other peripheral vasodilator particularly in patients with severe Aortic Stenosis.

SIDE EFFECT

Amlodipine is well tolerated. But Headache, Edema, Fatigue, Nausea, Flushing, Dizziness, Gum Hyperplasia and Erythema multiforme are reported.

USE IN PREGNANCY AND LACTATION

Pregnancy: Safety in pregnancy of Amlodipine has not been established.

Lactation: It is not known whether Amlodipine is excreted in breast milk. It is advised to stop breastfeeding during treatment with Amlodipine.

DRUG INTERACTION

Digoxin: Absence of any interaction between Amlodipine and Digoxin in healthy volunteers has been documented in a controlled clinical study Warfarin: A study showed that Amlodipine does not significantly alter the effect of

Warfarin on Prothrombin time. Cimetidine: A study showed that no interaction between Amlodipine & Cimetidine in

healthy volunteers.

PACKAGING

Lodical 5 tablet: Each box contains 3x10/5x10/6x10/10x10 tablets in blister pack.

